

KAWACHI'S INVARIANT FOR NORMAL SURFACE SINGULARITIES

VLADIMIR MAŠEK

ABSTRACT. We study a useful numerical invariant of normal surface singularities, introduced recently by T. Kawachi. Using this invariant, we give a quick proof of the (well-known) fact that all log-canonical surface singularities are either elliptic Gorenstein or rational (without assuming a priori that they are $\mathbb{Q}$ -Gorenstein).

In §2 we prove effective results (stated in terms of Kawachi's invariant) regarding global generation of adjoint linear systems on normal surfaces with boundary. Such results can be used in proving effective estimates for global generation on singular threefolds. The theorem of Ein–Lazarsfeld and Kawamata, which says that the minimal center of log-canonical singularities is always normal, explains why the results proved here are relevant in that situation.

Contents.

- 0. Introduction
- 1. Kawachi's invariant and log-canonical singularities
- 2. A theorem of Reider type on normal surfaces with boundary

Notations.

$\lceil \cdot \rceil$	round-up
$\lfloor \cdot \rfloor$	round-down
$\{\cdot\}$	fractional part
$f^{-1}D$	strict transform (proper transform)
f^*D	pull-back (total inverse image)
$\equiv$	numerical equivalence
$\sim$	linear equivalence

0. INTRODUCTION

Let Y be a normal algebraic surface over an algebraically closed field of arbitrary characteristic. Let $y \in Y$ be a fixed point on Y . Let $f : X \rightarrow (Y, y)$ be the minimal resolution of the germ (Y, y) if y is singular, resp. the blowing-up of Y at y if y is smooth. Kawachi ([8]) introduced the following numerical invariant of (Y, y) :

Definition. $\delta_y = -(Z - \Delta)^2$, where Z is the fundamental cycle of y and $\Delta = f^*K_Y - K_X$ is the canonical cycle (or antidiscrepancy) of y .

In §1 we recall several definitions (including the fundamental and the canonical cycle, etc.); then we study Kawachi's invariant and we give a very short proof

of the fact (well-known to the experts) that “numerically” log-canonical surface singularities are automatically $\mathbb{Q}$ -Gorenstein.

In §2 we prove several criteria for global generation of linear systems of the form $|K_Y + \lceil M \rceil|$, M a $\mathbb{Q}$ -divisor on Y such that $K_Y + \lceil M \rceil$ is Cartier. This type of results was the original motivation for introducing the invariant δ_y (see [8, 9, 10]). The main interest in results of Reider type for $\mathbb{Q}$ -divisors on normal surfaces comes from work related to Fujita’s conjecture for (log-) terminal threefolds, cf. [5, 11]. Using the criterion proved in §2, together with other recent results, we can significantly improve the main theorem of [5]; we will do so in a forthcoming paper.

The author has benefitted from numerous discussions with L. Bădescu, L. Ein, T. Kawachi, R. Lazarsfeld, and N. Mohan Kumar.

1. KAWACHI’S INVARIANT AND LOG-CANONICAL SINGULARITIES

In this section we recall several standard definitions and facts regarding normal surface singularities and we prove a number of elementary lemmas involving Kawachi’s invariant. For convenience, we use M. Reid’s recent notes [13] as our main reference.

(1.1) Let Y be a complete normal algebraic surface (in any characteristic), and let $f : X \rightarrow Y$ be a resolution of singularities of Y . We use Mumford’s $\mathbb{Q}$ -valued pullback and intersection theory on Y (cf. [12]): if D is any Weil (or $\mathbb{Q}$ -Weil) divisor on Y , then $f^*D = f^{-1}D + D_{\text{exc}}$, where $f^{-1}D$ is the strict transform of D and D_{exc} is the (unique) f -exceptional $\mathbb{Q}$ -divisor on X such that $f^*D \cdot F = 0$ for every f -exceptional curve $F \subset X$. (The existence and uniqueness of D_{exc} follow from the negative definiteness of the intersection form on exceptional curves.) If D_1, D_2 are two $\mathbb{Q}$ -Weil divisors on Y , then $D_1 \cdot D_2 \stackrel{\text{def}}{=} f^*D_1 \cdot f^*D_2$. See, e.g., [5, §1] for a quick review of this theory. [Here are a few key points: $D \geq 0 \implies f^*D \geq 0$; $D_1 \cdot D_2$ is independent of resolution; if C is a $\mathbb{Q}$ -divisor on X , then $C \cdot f^*D = f_*C \cdot D$; if D is $\mathbb{Q}$ -Cartier, then the definition of f^*D coincides with the usual one.]

Definition. (a) Let M be a $\mathbb{Q}$ -Weil divisor on Y . Then M is *nef* if $M \cdot C \geq 0$ for all irreducible curves $C \subset Y$. (Equivalently, M is nef if and only if f^*M is nef on the smooth surface X .)

(b) Assume that M is nef. Then M is *big* if in addition $M^2 > 0$ (i.e., if f^*M is big on X).

(1.2) Now let $y \in Y$ be a fixed point, and let $f : X \rightarrow (Y, y)$ be the *minimal* resolution of the germ (Y, y) if y is singular, resp. the blowing-up of Y at y if y is smooth. Let $f^{-1}(y) = \cup_{j=1}^N F_j$ (set-theoretically); $N = 1$ if y is smooth.

Definition. The *fundamental cycle* of (Y, y) is the smallest nonzero effective divisor $Z = \sum z_j F_j$ on X (with $z_j \in \mathbb{Z}$) such that $Z \cdot F_j \leq 0, \forall j$ (cf. [13, 4.5] or [2, p.132]).

Note that $z_j \geq 1, \forall j$, because $\cup F_j$ is connected.

If y is smooth, then $Z = F_1$ (F_1 is a (-1) -curve in this case).

Let $p_a(Z) = \frac{1}{2}Z \cdot (Z + K_X) + 1$; then $p_a(Z) \geq 0$, cf. [2, Theorem 3].

(1.3) Let K_X be a canonical divisor on X ; then f_*K_X is a canonical divisor K_Y on Y , and $\Delta \stackrel{\text{def}}{=} f^*K_Y - K_X$ is an f -exceptional $\mathbb{Q}$ -divisor on X , $\Delta = \sum a_j F_j$. Note that Δ is f -numerically equivalent to K_X ($\Delta \equiv_f -K_X$), i.e. $\Delta \cdot F_j = -K_X \cdot F_j, \forall j$; in particular, $p_a(Z) = \frac{1}{2}Z \cdot (Z - \Delta) + 1$, or $Z \cdot (Z - \Delta) = 2p_a(Z) - 2$.

If y is smooth, then $\Delta = -F_1$. On the other hand, if y is singular and f is the minimal resolution of (Y, y) , then $\Delta \cdot F_j = -K_X \cdot F_j \leq 0, \forall j$, so that Δ is effective (see, e.g., [5, Lemma 1.4]). In fact, $\Delta = 0$ if and only if y is a canonical singularity ($= RDP$, $= du\ Val$ singularity), cf. the last part of (1.4) below, and if y is not canonical, then $a_j > 0$ for all $j = 1, \dots, N$ (again, because $\cup F_j$ is connected).

Definition. Δ is the *canonical cycle* (or *antidiscrepancy*) of (Y, y) .

Δ is uniquely defined, even though K_X (and, accordingly, K_Y) is defined only up to linear equivalence. This follows from $\Delta \cdot F_j = -K_X \cdot F_j, \forall j$, and the negative definiteness of $\|F_i \cdot F_j\|$.

Assume that y is Gorenstein and non-canonical. Then Δ has integer coefficients and $\Delta > 0$; by the definition of the fundamental cycle, $\Delta \geq Z$. On the other hand, if (Y, y) is a log-canonical singularity, then $\Delta \leq Z$ (see below).

(1.4) Definition. y is a *rational* singularity if $R^1 f_* \mathcal{O}_X = 0$, or equivalently (cf. [2, Theorem 3]), if $p_a(Z) = 0$.

If y is any normal singularity and D is a Cartier divisor on Y , then f^*D has integer coefficients. If y is a rational singularity, then the converse is also true: if f^*D has integer coefficients, then D is Cartier. (*Proof:* $f^*D \cdot F_j = 0, \forall j$, and therefore f^*D is trivial in an open neighborhood of $f^{-1}(y)$, cf. [2, Lemma 5]; thus D is trivial in a punctured open neighborhood of y in Y , and therefore D is Cartier, because y is normal.) In particular, a rational surface singularity is always $\mathbb{Q}$ -factorial.

If y is a rational singularity, then $\text{mult}_y Y = -Z^2$ (cf. [2, Corollary 6], or [13, 4.17]); in particular, y is a *RDP* $\iff Z^2 = -2 \iff Z \cdot \Delta = 0 \iff \Delta \cdot F_j = 0, \forall j \iff \Delta = 0$. (Therefore *RDP* $\implies$ Gorenstein, because $f^*K_Y = K_X$ has integer coefficients, and therefore K_Y is Cartier. The converse is also true: rational Gorenstein $\implies$ *RDP*. Indeed, if y is Gorenstein, then either $\Delta = 0$ or $\Delta \geq Z$; but $\Delta \geq Z \implies Z \cdot (Z - \Delta) \geq 0 \implies p_a(Z) \geq 1$.)

(1.5) Definition. y is an *elliptic* singularity if $R^1 f_* \mathcal{O}_X$ is 1-dimensional. It is *elliptic Gorenstein* if in addition it is Gorenstein.

Lemma 1. y is *elliptic Gorenstein* if and only if $Z = \Delta$.

Proof. $\implies$ is proved in [13, 4.21]. Conversely, if $Z = \Delta$, then $K_X + Z = f^*K_Y$, so that $(K_X + Z) \cdot F_j = 0, \forall j$; the proof proceeds as in [13, 4.21] (go directly to Step 3 there). $\square$

(1.6) Definition. y is *log-terminal* (resp. *log-canonical*) if $a_j < 1$ (resp. $a_j \leq 1$), $\forall j$ – where $\Delta = f^*K_Y - K_X = \sum a_j F_j$, as before.

In dimension three or higher, one must assume that y is $\mathbb{Q}$ -Gorenstein before defining Δ (and log-terminal singularities, etc.) Using Mumford's definition of f^*K_Y , we don't need to make this assumption in the two-dimensional case. What's more, we will see in a moment that log-canonical (in our sense) automatically implies $\mathbb{Q}$ -Gorenstein. (This is also clear from the complete list of all log-canonical singularities: the arguments in [1] do not use the $\mathbb{Q}$ -Gorenstein condition.)

(1.7) Definition. $\delta_y = -(Z - \Delta)^2$.

Thus $\delta_y \in \mathbb{Q}$, $\delta_y \geq 0$, and $\delta_y = 0 \iff Z = \Delta \iff y$ is elliptic Gorenstein, by Lemma 1.

Lemma 2.

- (a) $\delta_y = 2 - 2p_a(Z) - \sum_{j=1}^N (z_j - a_j) K_X \cdot F_j$.
- (b) $\delta_y = 2 - 2p_a(Z) + \sum_{j=1}^N a_j (Z - \Delta) \cdot F_j$.

Proof. (a) $\delta_y = -(Z - \Delta)^2 = -Z \cdot (Z - \Delta) + \Delta \cdot (Z - \Delta) = 2 - 2p_a(Z) - K_X \cdot (Z - \Delta) = 2 - 2p_a(Z) - \sum_{j=1}^N (z_j - a_j) K_X \cdot F_j$ (because $\Delta \equiv_f K_X$).

(b) is similar. □

Lemma 3. (cf. [10, Theorem 1])

- (a) $\delta_y = 4$ if y is smooth;
- (b) $\delta_y = 2$ if y is a RDP;
- (c) $0 < \delta_y < 2$ if y is log-terminal but not smooth or a RDP;
- (d) $0 \leq \delta_y \leq 2$ if y is log-canonical but not smooth.

Proof. (a) y smooth $\implies Z - \Delta = 2F_1$, and $F_1^2 = -1$; thus $\delta_y = 4$.

(b) y a RDP $\implies \Delta = 0$, and $\delta_y = -Z^2 = \text{mult}_y Y = 2$.

(c), (d) δ_y is always ≥ 0 . If $\delta_y = 0$, then $\Delta = Z$, so that $a_j = z_j \geq 1, \forall j$; therefore y log-terminal $\implies \delta_y > 0$.

On the other hand, log-canonical $\implies a_j \leq 1 \leq z_j, \forall j$, so that $\delta_y \leq 2$ by Lemma 2 above ($K_X \cdot F_j \geq 0, \forall j$, because f is the *minimal* resolution).

Finally, if y is log-terminal, then $a_j < z_j, \forall j$; then $\delta_y = 2 \implies p_a(Z) = 0$ and $K_X \cdot F_j = 0, \forall j$ (recall that $p_a(Z) \geq 0$ for all normal singularities). Thus y is a rational singularity, and the last paragraph of (1.4) shows that y is a RDP. □

Corollary of the proof. *If y is log-terminal then it is rational. If y is log-canonical then it is either elliptic Gorenstein or rational. In particular, log-canonical implies $\mathbb{Q}$ -Gorenstein. (We noted already in (1.4) that rational implies $\mathbb{Q}$ -factorial.)*

Indeed, the proof above shows that y log-canonical $\implies p_a(Z) \leq 1$ (because $\delta_y \geq 0, z_j - a_j \geq 0$, and $K_X \cdot F_j \geq 0, \forall j$); moreover, $p_a(Z) = 1 \implies \delta_y = 0 \implies Z = \Delta \implies y$ is elliptic Gorenstein – and this can happen only in the log-canonical, not in the log-terminal case. In all other cases, $p_a(Z) = 0$, and therefore y is rational.

(1.8) The invariant $-\Delta^2$ has been considered before; for example, Sakai [15] proved results of Reid type on normal surfaces, using this invariant. In a sense, $-\Delta^2$ may be viewed as a local analogue of the Chern number c_1^2 .

At least in the rational case, $-\Delta^2$ is closely related to $\text{mult}_y Y$, via Kawachi's invariant:

Lemma 4. $-\Delta^2 = -Z^2 + \delta_y + 4(p_a(Z) - 1)$

Proof. $\Delta = Z - (Z - \Delta)$; therefore

$$-\Delta^2 = -Z^2 - (Z - \Delta)^2 + 2Z \cdot (Z - \Delta) = -Z^2 + \delta_y + 4(p_a(Z) - 1) \quad \square$$

Corollary 5. *If y is rational, then $p_a(Z) = 0$ and $-Z^2 = \text{mult}_y Y$, so that*

$$-\Delta^2 = \text{mult}_y Y - (4 - \delta_y).$$

In particular, if y is log-terminal but not canonical, we get:

$$\text{mult}_y Y - 4 < -\Delta^2 < \text{mult}_y Y - 2.$$


(1.9) Since log-canonical surface singularities are classified (see, for example, [1, 7]), one could compute δ_y explicitly in all cases. Indeed, this was Kawachi's original proof of part (c) in Lemma 3; see [8] for the complete list in the log-terminal case.

The computation of Z, Δ , and δ_y is an easy exercise in linear algebra. For illustration, we give here the values of Z, Δ , and δ_y for the “truly” log-canonical (i.e. non-log-terminal) singularities. The list of all such singularities can be found in [1, p.58]. To simplify notation, we assume that the chains of smooth rational curves shown in [1] consist of just one curve each. In each case, we show the dual graph of $\cup F_j$, indicating the self-intersection numbers F_j^2 .

Notice that in the “truly” log-canonical case δ_y takes only the values 0, 1, and 2 (always an integer), and the value of δ_y distinguishes three types of log-canonical singularities (with one interesting exception, noted below):


Notation: $\odot$ = smooth elliptic curve; $\circ$ = smooth rational curve.

Type 1: elliptic Gorenstein (cases (4) and (5) in [1])




$$Z = \Delta = F_1, \text{ resp. } F_1 + \cdots + F_n; \quad \delta_y = 0.$$

Type 2: (case (6) in [1])



$$\begin{aligned} w &\geq 2; \quad (a, b, c) = (3, 3, 3), (2, 2, 4), \text{ or } (2, 3, 6) \\ Z &= F_1 + (F_2 + F_3 + F_4) \text{ if } w \geq 3, Z = 2F_1 + (F_2 + F_3 + F_4) \text{ if } w = 2; \\ \Delta &= F_1 + (1 - \frac{1}{a})F_2 + (1 - \frac{1}{b})F_3 + (1 - \frac{1}{c})F_4; \quad \delta_y = 1. \end{aligned}$$

Type 3: (cases (7) – (8) in [1])




$w \geq 3$. If $w \geq 4$, then:

$$Z = F_1 + (F_2 + \cdots + F_5); \quad \Delta = F_1 + \frac{1}{2}(F_2 + \cdots + F_5); \quad \delta_y = 2.$$

However, if $w = 3$, then $Z = 2F_1 + (F_2 + \cdots + F_5)$ (while Δ is the same), and $\delta_y = 1$. This exceptional case illustrates an interesting property of the fundamental cycle; see (2.10) below.

(1.10) Exercise. Calculate Z , Δ , and δ_y for the following dual graph (cf. [3, p.350]):



(1.11) We conclude this section with another example. Assume that $f^{-1}(y)$ is a smooth curve C of genus g , with $C^2 = -w, w \geq 1$. This situation can be realized easily in practice: for example, y could be the vertex of the cone $Y = \text{Proj}(\bigoplus_{k \geq 0} H^0(C, kL))$, with C an arbitrary smooth curve of genus g and L an arbitrary divisor of degree w on C .

$$\text{Then } Z = C, \quad \Delta = \left(\frac{2}{w}(g-1) + 1\right)C, \quad \text{and} \quad \delta_y = \frac{4(g-1)^2}{w}.$$

If $g = 0$, then $\Delta = (1 - \frac{2}{w})C$, y is log-terminal, and $\delta_y = \frac{4}{w}$. If $w = 1$ then y is smooth. If $w = 2$ then y is an A_1 singularity (ordinary double point). If $w \geq 3$ then y is a log-terminal singularity “of type A_1 ”.

If $g = 1$, then y is log-canonical and $\delta_y = 0$. Such a singularity is known as *simply elliptic*.

If $g \geq 2$, then y is not log-canonical. Note that δ_y can be arbitrarily large in this case (if g is large relative to w). In particular, δ_y may be greater than 4 (which is the value for smooth points).

2. A THEOREM OF REIDER TYPE ON NORMAL SURFACES WITH BOUNDARY

In this section the ground field is $\mathbb{C}$.

(2.1) Let Y be a projective surface over $\mathbb{C}$, and let y be a fixed point on Y . Assume that Y is smooth except perhaps at y , which may be either smooth or a *RDP*. Let M be a nef and big $\mathbb{Q}$ -divisor on Y , with the property that $[M]$ is Cartier.

Ein and Lazarsfeld proved the following criterion on global generation:

Let $B = [M] - M$, and let $\mu = \text{mult}_y B$ if y is smooth, resp. $\mu = \max\{t \geq 0 \mid f^*B \geq tZ\}$ when y is a *RDP*, where $f : X \rightarrow (Y, y)$ is the minimal resolution and Z is the fundamental cycle.

Theorem. ([4, Theorem 2.3]) *Assume that $M^2 > (2 - \mu)^2$ and $M \cdot \mathbb{C} \geq (2 - \mu)$ for all curves C through y (when y is smooth), resp. that $M^2 > 2 \cdot (1 - \mu)^2$ and $M \cdot C \geq (1 - \mu)$ for all C through y (when y is a *RDP*).*

Then $y \notin \text{Bs}|K_Y + [M]|$.

This theorem was an important part of Ein and Lazarsfeld's proof of Fujita's conjecture on smooth threefolds. In extending that work to (log-) terminal threefolds (as required by the minimal model theory), it was necessary to extend the criterion mentioned above to arbitrary normal surfaces. Such extensions were obtained, e.g., in [5, Theorem 1.6], [11, Theorem 7]. However, these generalizations, while effective, are not optimal.

In a somewhat different direction, Kawachi and the author proved the following criterion, of independent interest:

Theorem. ([10, Theorem 2]) *Let Y be a normal surface, and let y be a fixed point on Y . Let $\delta = \delta_y$ if y is log-terminal, $\delta = 0$ otherwise. Let M be a nef divisor (with integer coefficients) on Y , such that $M^2 > \delta$ and $K_Y + M$ is Cartier.*

If $y \in \text{Bs}|K_Y + [M]|$, then there exists an effective divisor C passing through y , such that $M \cdot C < \frac{1}{2}\delta$ and $C^2 \geq M \cdot C - \frac{1}{4}\delta$ (in particular, y must be log-terminal, because M is nef).

Remark. When y is smooth, this is equivalent to Reider's original criterion ([14, Theorem 1]). When y is a *RDP*, we recover the Ein–Lazarsfeld criterion, plus a lower bound on C^2 .

While this result has several applications to linear systems on normal surfaces (cf. [10]), it cannot be used in the context of Fujita's conjecture on terminal threefolds, because M is required to have integer coefficients.

Kawachi formulated the following criterion for $\mathbb{Q}$ -divisors on normal surfaces:

Let Y, y be as before. Let $f : X \rightarrow (Y, y)$ be the minimal resolution if y is singular, resp. the blowing-up at y if y is smooth. Let Z and Δ be the fundamental and the canonical cycle, respectively. Let $\mu = \max\{t \geq 0 \mid f^*B \geq t(Z - \Delta)\}$; note that $\mu = 2 \cdot \text{mult}_y Y$ if y is smooth.

Open Problem. (cf. [9]) Let $\delta = \delta_y$ if y is log-terminal, $\delta = 0$ otherwise. Let M be a nef $\mathbb{Q}$ -Weil divisor on Y , such that $K_Y + B + M$ is Cartier, where $B = \lceil M \rceil - M$.

If $M^2 > (1 - \mu)^2 \delta$ and $M \cdot C \geq (1 - \mu) \frac{\delta}{2}$ for every curve C through y , then $y \notin \text{Bs}|K_Y + \lceil M \rceil|$.

When M has integer coefficients, this criterion is the same as the one mentioned above, minus the lower bound on C^2 . Also, this criterion contains the Ein–Lazarsfeld results for smooth and rational double points.

Kawachi formulated this as a theorem. Unfortunately his proof, based on a case-by-case analysis, is incomplete. In this section we prove a slightly weaker version, requiring that $M \cdot C \geq (1 - \mu)$, rather than $\geq (1 - \mu) \frac{\delta}{2}$, for all C through y . For application to Fujita’s conjecture on singular threefolds this makes little difference, though, because δ_y cannot be controlled in that situation anyway; the bound $\delta_y \leq 2$ (for y singular) is used instead.

(2.2) Let Y be a normal surface (= compact, normal two-dimensional algebraic space over $\mathbb{C}$). Let $y \in Y$ be a given point, and let $B = \sum b_i C_i$ be an effective $\mathbb{Q}$ -Weil divisor on Y with all $b_i \in \mathbb{Q}$, $0 \leq b_i \leq 1$; the C_i are distinct irreducible curves on Y . Since later we may need to consider more curves C_i than there are in $\text{Supp}(B)$, we allow some coefficients b_i to be 0. Let $f : X \rightarrow (Y, y)$ be the minimal resolution of singularities of the germ (Y, y) – resp. the blowing-up at y if y is a smooth point. Let $f^{-1}(y) = \cup F_j$, $Z = \sum z_j F_j$, $\Delta = \sum a_j F_j$, as in §1.

(2.3) Let $D_i = f^{-1}C_i$. Write $f^*B = f^{-1}B + B_{\text{exc}} = \sum b_i D_i + \sum b'_j F_j$.

Definition. (Y, B, y) is log-terminal (respectively log-canonical) if $a_j + b'_j < 1$ (respectively ≤ 1) for all j .

Thus (Y, y) is log-terminal (log-canonical) if $(Y, 0, y)$ is.

Remark. We do not require $K_Y + B$ to be $\mathbb{Q}$ -Cartier at y (unlike the similar definition in higher dimension); as in §1, this is a *consequence* of the other conditions. Note that $B \geq 0 \implies f^*B \geq 0$, and therefore (Y, B, y) log-terminal (log-canonical) $\implies (Y, y)$ log-terminal (log-canonical). Moreover, if $y \in \text{Supp}(B)$, then $b'_j > 0$, $\forall j$, and therefore (Y, B, y) log-canonical already implies (Y, y) log-terminal.

(2.4) Definition. Assume (Y, B, y) is log-terminal. Define

$$\mu = \mu_{B, y} = \max\{t \geq 0 \mid f^*B \geq t(Z - \Delta)\}.$$

Remark. All the z_j are ≥ 1 and all the a_j are < 1 ; μ is given explicitly by

$$\mu = \min \left\{ \frac{b'_j}{z_j - a_j} \right\}.$$

Of course, $\mu = 0$ if $B = 0$ (or, more generally, if $y \notin \text{Supp}(B)$). Note that $\mu = \frac{1}{2} \text{mult}_y(B)$ if y is a smooth point of Y .

Lemma 6. If (Y, B, y) is log-terminal and μ is defined as above, then $0 \leq \mu < 1$.

Indeed, $\mu \geq 0$ is clear. On the other hand, if $\mu \geq 1$ then we have $f^*B \geq (Z - \Delta)$, or $\Delta + f^*B \geq Z$, and therefore $a_j + b'_j \geq z_j \geq 1$ for every j , contradicting log-terminality. $\square$

(2.5) Let Y be a normal surface and $y \in Y$ a given point, as in (2.1). Let M be a **nef and big** $\mathbb{Q}$ -Weil divisor on Y such that $K_Y + \lceil M \rceil$ is Cartier. Let $B = \lceil M \rceil - M = \sum b_i C_i$; B is an effective $\mathbb{Q}$ -Weil divisor on Y , with $0 \leq b_i < 1$, $\forall i$.

We will prove the following criteria for freeness at y :

Theorem 7. *If (Y, B, y) is **not** log-terminal, then $y \notin \text{Bs } |K_Y + \lceil M \rceil|$.*

In the following two theorems, assume that (Y, B, y) is log-terminal. Then (Y, y) is also log-terminal. Define μ as in (2.3) and δ_y as in §1.

Also, assume that y is *singular*.

Theorem 8. *Assume that $M^2 > (1 - \mu)^2 \delta_y$ and $M \cdot C \geq (1 - \mu)$ for every curve $C \subset Y$ passing through y . (Note that M is still assumed to be nef, i.e. $M \cdot C \geq 0$ even for curves $C \subset Y$ not passing through y .)*

Then $y \notin \text{Bs } |K_Y + \lceil M \rceil|$.

In fact, if y is a singularity of type D_n or E_n (see [7, Remark 9.7]), we don't even need the assumption on $M \cdot C$ for C through y :

Theorem 9. *Assume that (Y, y) is a log-terminal singularity of type D_n or E_n , and M is nef and $M^2 > (1 - \mu)^2 \delta_y$. If (Y, y) is of type D_n , assume moreover that $M \cdot C > 0$ for every C through y .*

Then $y \notin \text{Bs } |K_Y + \lceil M \rceil|$.

(2.6) First we reduce the proof of Theorems 7, 8 and 9 to the case when y is the only singularity of Y :

Lemma 10. *We may assume that $Y - \{y\}$ is smooth.*

Proof. If y is not the only singularity of Y , then let $g : S \rightarrow Y$ be a simultaneous resolution of all singularities of Y *except* y . Put $M' = g^*M$ and $y' = g^{-1}(y)$ (note that g is an isomorphism of an open neighborhood of y' onto an open neighborhood of y). $K_S + \lceil M' \rceil$ is Cartier: outside y' this is clear, because $S - \{y'\}$ is smooth and $K_S + \lceil M' \rceil$ has integer coefficients, and in a certain open neighborhood of y' this is also clear, because g is an isomorphism there, and $K_Y + \lceil M \rceil$ is Cartier by hypothesis. Also, all the numerical conditions on M are satisfied by M' (in each of the hypotheses of Theorems 7, 8 and 9). If the theorems are true for S , y' , M' , then we get $y' \notin \text{Bs } |K_S + \lceil M' \rceil|$.

Write $\Delta' = g^*K_Y - K_S$; Δ' is an effective $\mathbb{Q}$ -divisor on S , and we have: $K_S + \lceil M' \rceil = \lceil g^*K_Y - \Delta' + g^*\lceil M \rceil - g^*B \rceil = g^*(K_Y + \lceil M \rceil) - \lfloor \Delta' + g^*B \rfloor$ (note that $K_Y + \lceil M \rceil$ is Cartier by hypothesis, and therefore $g^*(K_Y + \lceil M \rceil)$ has integer coefficients). Write $N = \lfloor \Delta' + g^*B \rfloor$; N is a divisor with integer coefficients on S , and $y' \notin \text{Supp}(N)$, because in a certain neighborhood of y' , Δ' is zero and g^*B is identified to B – whose coefficients are all < 1 .

In the first part of the proof we found a section $s \in H^0(S, g^*(K_Y + \lceil M \rceil) - N)$ which doesn't vanish at y' . Multiplying s by a global section of $\mathcal{O}_S(N)$ whose zero locus is N , we find a new section $t \in H^0(S, g^*(K_Y + \lceil M \rceil))$, which still doesn't vanish at y' . In turn, t corresponds to a global section of $\mathcal{O}_Y(K_Y + \lceil M \rceil)$ which doesn't vanish at y . $\square$

(2.7) Now assume that y is the only singularity of Y . Let $f : X \rightarrow Y$ be the minimal *global* desingularization of Y , $\Delta = f^*K_Y - K_X = \sum a_j F_j$, $B = \sum b_i C_i$, $f^*B = \sum b_i D_i + \sum b'_j F_j$, $Z = \sum z_j F_j$, etc.

Proof of Theorem 7

Assume that (Y, B, y) is not log-terminal; then there is at least one j such that $a_j + b'_j \geq 1$. f^*M is nef and big on X , and therefore the Kawamata–Viehweg vanishing theorem (cf. [4, Lemma 1.1]) gives: $H^1(X, K_X + \lceil f^*M \rceil) = 0$.

$K_X + \lceil f^*M \rceil = \lceil f^*K_Y - \Delta + f^*\lceil M \rceil - f^*B \rceil = f^*(K_Y + \lceil M \rceil) - \lfloor \Delta + f^*B \rfloor = f^*(K_Y + \lceil M \rceil) - \sum \lfloor a_j + b'_j \rfloor F_j = f^*(K_Y + \lceil M \rceil) - G$, where G is a nonzero effective divisor with integer coefficients on X such that $f(G) = \{y\}$. ($G > 0$ because $a_j + b'_j \geq 1$ for at least one j).

We have $H^1(X, f^*(K_Y + \lceil M \rceil) - G) = 0$, and therefore the restriction map

$$H^0(X, f^*(K_Y + \lceil M \rceil)) \rightarrow H^0(G, f^*(K_Y + \lceil M \rceil)|_G)$$

is surjective. As $f(G) = \{y\}$, $f^*(K_Y + \lceil M \rceil)|_G$ is trivial, i.e. it has a global section which doesn't vanish anywhere on G . By surjectivity, this section lifts to a global section $s \in H^0(X, f^*(K_Y + \lceil M \rceil))$ which doesn't vanish anywhere on G . In turn, s corresponds to a global section of $\mathcal{O}_Y(K_Y + \lceil M \rceil)$ which doesn't vanish at y , or else s would vanish everywhere on $f^{-1}(y)$, and in particular on G . $\square$

(2.8) Assume that y is log-terminal but not smooth. Thus $\Delta = \sum a_j F_j$ with $0 \leq a_j < 1$ for every j .

Proof of Theorem 8

Lemma 11. *If $M^2 > (1 - \mu)^2 \delta_y$, then we can find an effective $\mathbb{Q}$ -Weil divisor D on Y such that $D \equiv M$ and $f^*D \geq (1 - \mu)(Z - \Delta)$.*

Proof. Since $M^2 > (1 - \mu)^2 \delta_y$, $f^*M - (1 - \mu)(Z - \Delta)$ is in the positive cone of X (see [10], (2.3) for a similar argument). In particular, $f^*M - (1 - \mu)(Z - \Delta)$ is big.

Let $G \in |k(f^*M - (1 - \mu)(Z - \Delta))|$ for some k sufficiently large and divisible. Let $T = \frac{1}{k}G + (1 - \mu)(Z - \Delta)$. Write $T = \sum d_i D_i + \sum t_j F_j$. Define $D = f_*T = \sum d_i C_i$, and write $f^*D = \sum d_i D_i + \sum d'_j F_j$. We have:

1. $T \equiv f^*M$, because $kT \sim k f^*M$;
2. $T \geq 0$, and therefore $D \geq 0$;
3. $T \cdot F_j = 0$ for every exceptional curve F_j , because $T \equiv f^*M$;
4. $f^*D = T$; indeed, the coefficients d'_j are uniquely determined by the condition $f^*D \cdot F_j = 0$ for every j , and T already satisfies this condition;
5. $f^*D = T \geq (1 - \mu)(Z - \Delta)$, because $G \geq 0$;
6. finally, $D \equiv M$, because $f^*D = T \equiv f^*M$.

$\square$

Remark. $f^*D \geq (1 - \mu)(Z - \Delta)$ means $d'_j \geq (1 - \mu)(z_j - a_j)$ for every j . We may assume, however, that $d'_j > (1 - \mu)(z_j - a_j)$ for every j . Indeed, since $M^2 > (1 - \mu)^2 \delta_y$, we have $M^2 > (1 - \mu)^2 \delta_y (1 + \epsilon)^2$ for some small rational number $\epsilon > 0$; then working as above we can find $D \equiv M$ such that $d'_j \geq (1 - \mu)(z_j - a_j)(1 + \epsilon) > (1 - \mu)(z_j - a_j)$ for every j .

We'll assume that $d'_j > (1 - \mu)(z_j - a_j)$ for all j .

For every rational number c , $0 < c < 1$, let $R_c = f^*(M - cD)$. $R_c \equiv (1 - c)f^*M$, so that R_c is nef and big, and we have

$$(1) \quad H^1(X, K_X + \lceil R_c \rceil) = 0.$$

$$\begin{aligned} K_X + \lceil R_c \rceil &= \lceil f^*K_Y - \Delta + f^*\lceil M \rceil - f^*B - cf^*D \rceil \\ &= f^*(K_Y + \lceil M \rceil) - \lfloor \Delta + f^*B + cf^*D \rfloor \\ &= f^*(K_Y + \lceil M \rceil) - \sum [b_i + cd_i]D_i - \sum [a_j + b'_j + cd'_j]F_j. \end{aligned}$$

Choose $c = \min \left\{ \frac{1-a_j-b'_j}{d'_j}, \text{ all } j; \frac{1-b_i}{d_i}, \text{ all } i \text{ such that } d_i > 0 \text{ and } y \in C_i \right\}$.

Note that $c > 0$, because (Y, B, y) is log-terminal, and $c < 1$, because for every j we have $b'_j \geq \mu(z_j - a_j)$ and $z_j \geq 1$, and therefore $1 - a_j - b'_j \leq z_j - a_j - \mu(z_j - a_j) = (1 - \mu)(z_j - a_j) < d'_j$. Therefore (1) holds for this choice of c .

Note also that $0 < a_j + b'_j + cd'_j \leq 1$ for all j ; so $F \stackrel{\text{def}}{=} \sum [a_j + b'_j + cd'_j]F_j$ is either zero or a sum of distinct irreducible components, $F = F_1 + \dots + F_s$ (after re-indexing the F_j if necessary). Similarly, $\sum [b_i + cd_i]D_i = N + A$, where $\text{Supp}(N) \cap f^{-1}(y) = \emptyset$, and A is either zero or a sum of distinct irreducible components, $A = D_1 + \dots + D_t$, where $D_1, \dots, D_t$ meet $f^{-1}(y)$. Each component F_j of F (if any), and each component D_i of A (if any), has coefficient 1 in $\Delta + f^*B + cf^*D$. Also, F and A cannot both be equal to zero.

We will use the following form of the Kawamata–Viehweg vanishing theorem (see, for example, [4], Lemma 2.4):

Lemma 12. *Let X be a smooth projective surface over $\mathbb{C}$, and let R be a nef and big $\mathbb{Q}$ -divisor on X . Let $E_1, \dots, E_m$ be distinct irreducible curves such that $\lceil R \rceil \cdot E_i > 0$ for every i . Then*

$$H^1(X, K_X + \lceil R \rceil + E_1 + \dots + E_m) = 0.$$

We consider two cases, according to whether $F \neq 0$ or $F = 0$.

Case I: $F \neq 0$.

Using Lemma 12 for R_c in place of R and $D_1, \dots, D_t$ in place of $E_1, \dots, E_m$, we get $H^1(X, K_X + \lceil R_c \rceil + A) = 0$, or $H^1(X, f^*(K_Y + \lceil M \rceil) - N - F) = 0$. We conclude as in the proof of Theorem 7; the “ $-N$ ” part is treated as in the proof of Lemma 10. *Note:* $R_c \cdot D_i = (1 - c)M \cdot C_i > 0$ for every i , and D_i has integer coefficient in R_c if D_i is a component of A – and therefore $\lceil R_c \rceil \cdot D_i > 0$ for such D_i .

Case II: $F = 0$.

As noted earlier, in this case $A \neq 0$; using Lemma 12 as in Case I above, we get $H^1(X, f^*(K_Y + \lceil M \rceil) - N - D_1) = 0$, and therefore the restriction map

$$(2) \quad H^0(X, f^*(K_Y + \lceil M \rceil) - N) \rightarrow H^0(D_1, (f^*(K_Y + \lceil M \rceil) - N)|_{D_1})$$

is surjective.

$D_1 \cap f^{-1}(y) \neq \emptyset$; let $x \in D_1 \cap f^{-1}(y)$. Assume we can find a section $s' \in H^0(D_1, f^*(K_Y + \lceil M \rceil) - N|_{D_1})$ such that $s'(x) \neq 0$. Then by the surjectivity of (2) we can find $s \in H^0(X, f^*(K_Y + \lceil M \rceil) - N)$ such that $s(x) \neq 0$; then we conclude as in the proof of Lemma 10.

Hence the proof is complete if we show that $x \notin \text{Bs} |f^*(K_Y + \lceil M \rceil) - N|_{D_1}|$. Note that $f^*(K_Y + \lceil M \rceil) - N|_{D_1} = K_X + \lceil R_c \rceil + A|_{D_1} = (K_X + D_1)|_{D_1} + (\lceil R_c \rceil + D_2 + \cdots + D_t)|_{D_1} = K_{D_1} + (\lceil R_c \rceil + D_2 + \cdots + D_t)|_{D_1}$. By [6], Theorem 1.4 and Proposition 1.5, it suffices to show that $(\lceil R_c \rceil + D_2 + \cdots + D_t) \cdot D_1 > 1$ (then this intersection number is ≥ 2 , because it is an integer).

$\lceil R_c \rceil = R_c + \sum \{b_i + cd_i\}D_i + \sum \{a_j + b'_j + cd'_j\}F_j$. Note that $b_1 + cd_1 = 1$, and therefore $\{b_1 + cd_1\} = 0$. Also, since $F = 0$, we have $0 \leq a_j + b'_j + cd'_j < 1$ for every j – and therefore $\{a_j + b'_j + cd'_j\} = a_j + b'_j + cd'_j$. Hence we have

$$(\lceil R_c \rceil + D_2 + \cdots + D_t) \cdot D_1 \geq R_c \cdot D_1 + \sum (a_j + b'_j + cd'_j)F_j \cdot D_1.$$

$R_c \equiv (1 - c)f^*M$, so that $R_c \cdot D_1 = (1 - c)M \cdot C_1 \geq (1 - c)(1 - \mu)$, because $y \in C_1 = f_*D_1$. Also, D_1 meets at least one F_j , say F_1 . Therefore we have


$$\begin{aligned} (\lceil R_c \rceil + D_2 + \cdots + D_t) \cdot D_1 &\geq (1 - c)(1 - \mu) + (a_1 + b'_1 + cd'_1) \\ &> (1 - c)(1 - \mu) + a_1 + \mu(z_1 - a_1) + c(1 - \mu)(z_1 - a_1) \\ &\geq (1 - c)(1 - \mu) + a_1 + \mu(1 - a_1) + c(1 - \mu)(1 - a_1) \\ &= 1 + (1 - c)(1 - \mu)a_1 \\ &\geq 1. \end{aligned}$$

□

(2.9) Finally, we prove Theorem 9. We assume the reader is familiar with the classification of RDP 's. The classification of log-terminal singularities is similar: if $f : X \rightarrow (Y, y)$ is the minimal resolution of a log-terminal germ and $f^{-1}(y) = \cup F_j$, then the F_j are smooth rational curves, and the dual graph is a graph of type A_n , D_n or E_n (see, e.g., [7, §9], or [1]). The only difference is that in the log-terminal case the self-intersection numbers $-w_j = F_j^2$ are not necessarily all equal to -2 .

The classification of log-terminal singularities with non-zero reduced boundary is even simpler. We have:


Lemma 13. *Assume (Y, y) is a normal surface germ, and C_1 is a reduced, irreducible curve on Y such that $y \in C_1$. If (Y, C_1, y) is log-terminal, then (Y, y) is of type A_n . More explicitly, if $f : X \rightarrow (Y, y)$ is the minimal resolution, $f^{-1}(y) = F_1 \cup \cdots \cup F_n$, $-w_j = F_j^2$, and $f^{-1}C_1 = D_1$, then the dual graph of the resolution is:*



(If y is smooth, then $n = 1$ and $w_1 = 1$; otherwise all $w_j \geq 2$.)

Similarly, for log-canonical singularities with boundary we have:

Lemma 14. *Let (Y, y) and C_1 be as in the previous lemma. If (Y, C_1, y) is log-canonical, then either (Y, y) is of type A_n and the dual graph is the one shown above, or (Y, y) is of type D_n and the dual graph of the resolution is:*



All these facts can be found, for example, in [1], and also in [7].

Proof of Theorem 9

Going back to the proof of theorem 8, we will show that Case II of the proof is not possible if y is of type D_n or E_n . Therefore the condition $M \cdot C \geq (1 - \mu)$ for C through y is no longer necessary. We still need $M \cdot C > 0$ for C through y if $A \neq 0$, to be able to use Lemma 12 (see the *Note* at the end of Case I). However, we will show that $A = 0$ if y is of type E_n , so that in that case we don't even need the condition $M \cdot C > 0$ for C through y .

Assume that $A \neq 0$. Let C_1 be a component of A , and let $f^*C_1 = D_1 + \sum c'_j F_j$. Since C_1 is a component of A , we have $b_1 + cd_1 = 1$, and therefore $B + cD = C_1 + \text{other terms}$; consequently $f^*B + cf^*D \geq f^*C_1$, and in particular $b'_j + cd'_j \geq c'_j, \forall j$.

If we end up in Case II in the proof of Theorem 8, then $a_j + b'_j + cd'_j < 1$ for all j ; consequently $a_j + c'_j < 1, \forall j$, so that (Y, C_1, y) is log-terminal. By Lemma 13, (Y, y) must be of type A_n .

Now assume that we end up in Case I, with $A \neq 0$. We still have $a_j + b'_j + cd'_j \leq 1, \forall j$, and therefore (Y, C_1, y) is log-canonical. By Lemma 14, (Y, y) is either of type A_n or of type D_n . $\square$

Remark. For singularities of type D_n and E_n , Theorem 9 is stronger than the Open Problem (see (2.1)). To complete the proof of the Open Problem, the only case to consider is that of a singularity of type A_n .

In the proof of Theorem 8, Case II (which is possible only when y is of type A_n), we used the inequality $M \cdot C \geq (1 - \mu)$ for $y \in C$. In fact, using Lemma 13 and modifying slightly the final computation in (2.8), we see that we need slightly less: $M \cdot C \geq (1 - \mu)(1 - a)$, where $a = \min\{a_1, a_n\}$ (note that in Lemma 13, D_1 could meet either F_1 or F_n).

On the other hand, for y of type A_n we have $\delta_y = 2 - (a_1 + a_n)$ (for $n = 1$ this follows from (1.11); for $n \geq 2$ use Lemma 2, (b), and the obvious formulae $(Z - \Delta) \cdot F_j = -1$ for $j = 1$ and $j = n$, $(Z - \Delta) \cdot F_j = 0$ otherwise). Thus the Open Problem requires that $M \cdot C \geq (1 - \mu)(1 - \frac{a_1 + a_n}{2})$ for $y \in C$. In particular, the Open Problem is proved if $a_1 = a_n$ (e.g., if y is of type A_1).

(2.10) Analyzing the proofs of Theorems 8 and 9, we may ask: what was the relevance of Z being the fundamental cycle of y ? Δ arises naturally, as $f^*K_Y - K_X$; but Z could have been any effective, f -exceptional cycle with integer coefficients such that $z_j \geq 1$ for all j (i.e., such that $\text{Supp}(Z) = f^{-1}(y)$). The answer is provided by the following proposition:

Proposition 15. *Let (Y, y) be a log-terminal singularity, with $f : X \rightarrow (Y, y)$ the minimal resolution (resp. the blowing-up at y if y is smooth). Let Z and Δ be the*

fundamental, resp. the canonical cycle. Let $Z' = \sum z'_j F_j$ be any other effective, f -exceptional cycle (with integer coefficients), such that $z'_j \geq 1$ for all j .

Then $\delta_y \leq \delta'$, where $\delta_y = -(Z - \Delta)^2$ and $\delta' = -(Z' - \Delta)^2$. Moreover, Z is the (unique) largest cycle among all the Z' for which $\delta' = \delta_y$.

The proof depends on the detailed classification of log-terminal surface singularities, cf. [7]. Explicitly, we need the following lemma, which can be proved by brute force (the computations are straightforward in all cases; for reference, they can be found in [8]):

Lemma 16. *Let (Y, y) be log-terminal.*

(a) *There is at most one F_j with $(Z - \Delta) \cdot F_j > 0$. If such an F_j exists, then $(Z - \Delta) \cdot F_j = 1$ and the corresponding $w_j = -F_j^2 \geq 3$.*

(b) *There is at most one F_j with $z_j \geq 2$ and $(Z - \Delta) \cdot F_j < 0$. If such an F_j exists, then $z_j = 2$ and $(Z - \Delta) \cdot F_j = -1$.*

Proof of Proposition 15

Let $Z' = Z + (P - N)$ with $P, N \geq 0$ without common components. Then

$$\begin{aligned} \delta' &= -(Z' - \Delta)^2 = -(Z + P - N - \Delta)^2 \\ &= -(Z - \Delta)^2 - (P - N)^2 - 2(Z - \Delta) \cdot (P - N) \\ &= \delta_y + (-P^2) - 2(Z - \Delta) \cdot P + (-N^2) + 2(Z - \Delta) \cdot N + 2(P \cdot N). \end{aligned}$$

Since $P \cdot N \geq 0$, the Proposition is proved if we can prove that

- (a) $(-P^2) - 2(Z - \Delta) \cdot P > 0$ if $P > 0$;
- (b) $(-N^2) + 2(Z - \Delta) \cdot N \geq 0$.

Proof of (a).

If $P = \sum t_j F_j$, then we have

$$(-P^2) - 2(Z - \Delta) \cdot P = (-P^2) - 2 \sum t_j (Z - \Delta) \cdot F_j.$$

If $(Z - \Delta) \cdot F_j \leq 0$ for all j , then we are done (note that $(-P^2) > 0$ if $P > 0$). Otherwise there is exactly one j , call it j_0 , such that $(Z - \Delta) \cdot F_{j_0} > 0$. By Lemma 16, we have $w_{j_0} = -F_{j_0}^2 \geq 3$ and $(Z - \Delta) \cdot F_{j_0} = 1$. Therefore

$$(-P^2) - 2 \sum t_j (Z - \Delta) \cdot F_j \geq (-P^2) - 2t_{j_0}.$$

We will show that $(-P^2) \geq t_{j_0}^2 + 2$; then (a) will follow.

Write $F_j^2 = -w_j$, $F_i \cdot F_j = l_{ij}$ for $i < j$ ($l_{ij} = 1$ if F_i meets F_j , 0 otherwise). We have:

$$\begin{aligned} (-P^2) &= \sum_j w_j t_j^2 - \sum_{i < j} 2l_{ij} t_i t_j \\ &\geq t_{j_0}^2 + \sum_j 2t_j^2 - \sum_{i < j} 2l_{ij} t_i t_j \end{aligned}$$

(note that $w_j \geq 2$, $\forall j$, and $w_{j_0} \geq 3$).

Now consider a singularity (Y', y') , whose minimal resolution has a dual graph identical to that of (Y, y) , except that $F_j'^2 = -2$, $\forall j$ (thus y' is a “true” A_n, D_n ,

or E_n rational double point). If $P' = \sum t_j F'_j$ (having the same coefficients as P), then $(-P'^2) > 0$; that is,

$$\sum_j 2t_j^2 - \sum_{i < j} 2l_{ij}t_it_j > 0$$

— and therefore ≥ 2 , because it is an *even* integer. $\square$

Proof of (b).

If $N = \sum x_j F_j$, then we have


$$(-N^2) + 2(Z - \Delta) \cdot N = (-N^2) + 2 \sum x_j (Z - \Delta) \cdot F_j.$$

If $(Z - \Delta) \cdot F_j \geq 0$ for all j , or if $x_j = 0$ whenever $(Z - \Delta) \cdot F_j < 0$, then we are done. Note that $x_j = z_j - z'_j$ if $z_j > z'_j$, and 0 otherwise. Thus $x_j \geq 1 \implies z_j \geq 2$. Therefore, by Lemma 16, there can be at most one negative term in $2 \sum x_j (Z - \Delta) \cdot F_j$; and if there is one, corresponding, say, to j_1 , then $z_{j_1} = 2$ and $(Z - \Delta) \cdot F_{j_1} = -1$. Therefore $x_{j_1} = 1$, and $2(Z - \Delta) \cdot N \geq -2$. Finally, $(-N^2) \geq 2$ (if $N \neq 0$), as in the proof of (a) above. $\square$

Remarks.

1. We showed that $[-(Z' - \Delta)^2 = \delta_y] \implies [Z' \leq Z]$. The converse is not always true. For example, if F_j is a component with $z_j = 2$ and $(Z - \Delta) \cdot F_j = 0$ (such components exist in some cases of type D_n and E_n , cf. [8]), then taking $Z' = Z - F_j$ we get $-(Z' - \Delta)^2 > \delta_y$ (cf. the proof of (b) above).

2. If (Y, y) is not log-terminal, then the statement of Proposition 15 is no longer necessarily true, even if y is rational. For example:



$Z = F_1 + (F_2 + \cdots + F_6)$, $\Delta = \frac{6}{5}F_1 + \frac{3}{5}(F_2 + \cdots + F_6)$, and $\delta_y = \frac{13}{5}$; but $\delta' = -(Z' - \Delta)^2 = \frac{8}{5} < \delta_y$ for $Z' = Z + F_1$.

Exercise. Is the statement of Proposition 15 true for log-canonical singularities?

Hint: There is nothing to prove if $\delta_y \leq 1$. Notice that in the exceptional case of (1.9), Type 3 with $w = 3$, we have $\delta' = 2$ for $Z' = F_1 + (F_2 + \cdots + F_5)$, while $\delta_y = 1$ (in that case we have $Z = 2F_1 + (F_2 + \cdots + F_5)$).

REFERENCES

- [1] V. Alexeev, *Classification of log-canonical surface singularities*, in: *Flips and abundance for algebraic threefolds – Salt Lake City, Utah, August 1991*, J. Kollár (editor), Astérisque No. **211** (1992), Société Mathématique de France, 1992, 47–58.
- [2] M. Artin, *On isolated rational singularities of surfaces*, Amer. J. Math. **88** (1966), 129–136.
- [3] E. Brieskorn, *Rationale Singularitäten komplexer Flächen*, Invent. Math. **4** (1968), 336–358.

- [4] L. Ein and R. Lazarsfeld, *Global generation of pluricanonical and adjoint linear series on smooth projective threefolds*, J. Amer. Math. Soc. **6** (1993), 875–903.
- [5] L. Ein, R. Lazarsfeld and V. Mašek, *Global generation of linear series on terminal threefolds*, Internat. J. Math. **6** (1995), 1–18.
- [6] R. Hartshorne, *Generalized divisors on Gorenstein curves and a theorem of Noether*, J. Math. Kyoto Univ. **26** (1986), 375–386.
- [7] Y. Kawamata, *Crepant blowing-up of 3-dimensional canonical singularities and its applications to degeneration of surfaces*, Ann. Math. **127** (1988), 93–163.
- [8] T. Kawachi, *On freeness theorem of the adjoint bundle on a normal surface*, Duke e-print [alg-geom/9603022](#) (March 1996).
- [9] T. Kawachi, *Effective base point freeness on a normal surface*, Duke e-print [alg-geom/9612018](#) (December 1996).
- [10] T. Kawachi and V. Mašek, *Reider-type theorems on normal surfaces*, to appear in J. Alg. Geom.
- [11] D. Matsushita, *Effective base point freeness*, Kodai Math. J. **19** (1996), 87–116.
- [12] D. Mumford, *The topology of normal singularities of an algebraic surface and a criterion for simplicity*, Inst. Hautes Études Sci. Publ. Math. **9** (1961), 5–22.
- [13] M. Reid, *Chapters on Algebraic Surfaces*, in: *Complex Algebraic Geometry*, J. Kollár (editor), IAS/Park City Mathematics Series, Vol. **3** (1997), 1–160.
- [14] I. Reider, *Vector bundles of rank 2 and linear systems on algebraic surfaces*, Ann. Math. **127** (1988), 309–316.
- [15] F. Sakai, *Reider–Serrano’s method on normal surfaces*, Algebraic Geometry: Proceedings, L’Aquila 1988, Lect. Notes in Math., vol. 1417, 1990, 301–319.

DEPARTMENT OF MATHEMATICS, BOX 1146, WASHINGTON UNIVERSITY, ST. LOUIS, MO 63130
E-mail address: vmasek@math.wustl.edu